

On the validity of constraints on light elementary particles and extra-dimensional physics from the Casimir effect

E. Fischbach¹, G.L. Klimchitskaya^{2,3}, D.E. Krause^{4,1}, V.M. Mostepanenko^{5,3,a}

¹ Department of Physics, Purdue University, West Lafayette, Indiana 47907, USA

² North-West Technical University, Millionnaya St. 5, St.Petersburg, 191065, Russia

³ Institute for Theoretical Physics, Leipzig University, Postfach 100920, D-04009, Leipzig, Germany

⁴ Physics Department, Wabash College, Crawfordsville, Indiana 47933, USA

⁵ Noncommercial Partnership “Scientific Instruments”, Tverskaya St. 11, Moscow, 103905, Russia

Received: March 5, 2010 / Revised version: date

Abstract. We discuss the constraints on the parameters of a Yukawa interaction obtained from the indirect measurements of the Casimir pressure between parallel plates using the sphere-plate configuration. Recently, it was claimed in the literature that the application of the proximity force approximation (PFA) to the calculation of a Yukawa interaction in the sphere-plate configuration could lead to a large error of order 100% in the constraints obtained. Here we re-calculate the constraints both exactly and using the PFA, and arrive at identical results. We elucidate the reasons why an incorrect conclusion was obtained suggesting that the PFA is inapplicable to calculate the Yukawa force.

The Yukawa-type corrections to Newtonian gravitational law are predicted from the exchange of light elementary particles and in some extra-dimensional generalizations of the standard model. Within a short interaction range of around $0.1\,\mu\text{m}$ the strongest constraints on the parameters of Yukawa interaction were obtained from the indirect dynamic measurement of the Casimir pressure between two parallel plates using the configuration of an Au-coated sphere above an Au-coated plate of a micro-machined oscillator [1,2]. In [1,2] this was done using what is referred to as *the proximity force approximation* (PFA) [3] for the calculation of the Yukawa force acting between a sphere and a plate. For two point-like particles with masses m_1 and m_2 spaced at a separation r apart, the Yukawa corrections to Newtonian gravity are conventionally presented in the form [4,5]

$$V_{\text{Yu}}(r) = -\frac{\alpha G m_1 m_2}{r} e^{-r/\lambda}, \quad (1)$$

where G is the gravitational constant, α and λ are the strength and the interaction range of the Yukawa interaction, respectively.

Reference [6] especially investigated the possibility of calculating the gravitational and Yukawa-type interactions in sphere-plate configuration using the PFA. In so doing two formulations of the PFA were presented. In the most general formulation [7] the force of interaction is found as an integral of the known force between parallel surface elements. According to the simplified formulation of the

PFA, valid under certain conditions, the force is equal to $2\pi R E$ where R is the sphere radius and E is the energy per unit area in the configuration of two parallel plates [3]. In [6] the applicability of different formulations of the PFA for the calculation of Yukawa and gravitational interactions in the experimental configuration of [1,2] was confirmed. Specifically, it was shown that for the gravitational and Yukawa-type interactions between a compact body and a plane plate of infinitely large area the most general formulation of the PFA leads to the exact results. The simplified formulation of the PFA was shown to be approximately applicable to the Yukawa interaction.

Motivated by [6], the subsequent paper [8] discussed the same subject. In agreement with [6], paper [8] arrives at the conclusion that the level of precision in using the two formulations of the PFA to evaluate Yukawa forces in the sphere-plane geometry “is of the same order of magnitude as the Casimir theory-experiment comparison that uses PFA to compute the sphere-plate Casimir force.” On the basis of [9,10,11,12,13,14] it was also concluded [6] that for the experimental configuration of [1,2] with $a/R \approx 0.001$ (a is the closest separation between a sphere and a plate, R is the sphere radius) the relative error in the Casimir force due to the use of the PFA is about 0.1%. The same conclusion was reached in [8] on the basis of [11,12,15,16]. Specifically, [8] states that “the PFA... is expected to approximate the exact Casimir force within 0.1%.” Thus, in both papers [6,8] it is recognized that in recent experiment [1,2] the error in the evaluation of the Yukawa-type force between a sphere and a plate using the simplified formulation of the PFA is of about 0.1%.

^a E-mail: Vladimir.Mostepanenko@itp.uni-leipzig.de

However, notwithstanding the previous discussion, which supports the use of the PFA, [8] claims that the constraints on α , λ from the precision dynamic measurements of the Casimir force obtained in [1,2] could have “a large order of 100% correction.” According to [8], “Considering the relatively small margins of improvement reported recently (see for instance Fig. 3 in [21]), a systematic shift due to the use of the PFA instead of EPFA may lead to significant changes for the exclusion region in the $\alpha - \lambda$ plane.” We recall that [21] cited in [8] is just our reference [2] and EPFA is equivalent to the exact calculation. By way of contrast, [6] confirmed the validity of the constraints obtained in [1,2]. Since these constraints are now included in the Review of Particle Physics (Particle Data Group) for the year 2008 [17], it seems necessary to verify whether the claim of [8] which casts doubts on these constraints is correct. In what follows we repeat the derivation of the constraints on (α, λ) from the experimental data of [1,2] using the exact formulas for the Yukawa-type interaction, and arrive at the same results as obtained in [1,2] using the simplified formulation of the PFA. We show that the use of the simplified formulation of the PFA within its applicability region for the calculation of Yukawa-type forces cannot lead to any systematic shifts in the $\alpha - \lambda$ plane not only in already performed experiments but in presumably much more precise future experiments as well. Because of this it is not correct to call this formulation of the PFA “an invalid approximation to compute volumetric forces” [8]. As for any approximation, the simplified PFA is applicable in some ranges of parameters (see below) and is not applicable outside of these ranges. Several other misleading statements of [8] are also commented upon below.

To begin our derivation, it is pertinent to recall that in the experiment of [1,2] the separation distance between a sphere of $R = 151.3 \mu\text{m}$ radius and a plate was varied harmonically with time, and the directly measured quantity was the frequency shift of the oscillator under the influence of the Casimir force acting between the two bodies. Due to the properties of the oscillator, this frequency shift is proportional to the gradient of the Casimir force. Using the simplified PFA, this gradient turns out to be proportional to the effective Casimir pressure in the configuration of two parallel plates. Thus, in this experiment one indirectly measures the Casimir pressure between two parallel plates (see the recent review [18] for a detailed justification of this statement). To obtain constraints on the parameters of Yukawa-type interaction α and λ in [1,2,19] it was supposed that the gradient of the Yukawa-type force is proportional to the Yukawa-type pressure between the two parallel plates, i.e., the simplified PFA was applied to the Yukawa-type forces. As a result, the constraints were obtained from the inequality

$$|P^{\text{Yu}}(a)| \leq \tilde{\Xi}(a), \quad (2)$$

where $[-\tilde{\Xi}(a), \tilde{\Xi}(a)]$ is the minimum confidence interval [1] containing all differences $P^{\text{theor}}(a) - \bar{P}^{\text{expt}}(a)$ within the separation region $180 \text{ nm} < a < 746 \text{ nm}$. The Yukawa pressure between two parallel plates is given by

$$P^{\text{Yu}}(a) = -2\pi G\alpha\lambda^2 e^{-a/\lambda}$$

$$\begin{aligned} & \times \left[\rho_{\text{Au}} - (\rho_{\text{Au}} - \rho_{\text{Cr}}) e^{-\Delta_{\text{Au}}^{(s)}/\lambda} \right. \\ & \quad \left. - (\rho_{\text{Cr}} - \rho_s) e^{-(\Delta_{\text{Au}}^{(s)} + \Delta_{\text{Cr}}^{(s)})/\lambda} \right] \\ & \times \left[\rho_{\text{Au}} - (\rho_{\text{Au}} - \rho_{\text{Cr}}) e^{-\Delta_{\text{Au}}^{(p)}/\lambda} \right. \\ & \quad \left. - (\rho_{\text{Cr}} - \rho_{\text{Si}}) e^{-(\Delta_{\text{Au}}^{(p)} + \Delta_{\text{Cr}}^{(p)})/\lambda} \right]. \quad (3) \end{aligned}$$

Here, $\Delta_{\text{Au}}^{(s)}$, $\Delta_{\text{Cr}}^{(s)}$, $\Delta_{\text{Au}}^{(p)}$, and $\Delta_{\text{Cr}}^{(p)}$ are the thicknesses of the Au and Cr layers with the densities ρ_{Au} and ρ_{Cr} on the sphere and the plate, made of sapphire with density ρ_s and Si with density ρ_{Si} , respectively (the values of all parameters are contained in [1,2]).

To verify the constraints obtained in [1,2] from Eq. (2) based on the use of the simplified PFA for Yukawa forces, we return instead to the original inequality following from the experimental data, i.e., that the gradient of the Yukawa force normalized to $2\pi R$ belongs to the confidence interval determined at a 95% confidence level

$$\left| \frac{1}{2\pi R} \frac{\partial F_{sp}^{\text{Yu}}(a)}{\partial a} \right| \leq \tilde{\Xi}(a). \quad (4)$$

Here, $F_{sp}^{\text{Yu}}(a)$ is the Yukawa-type force for the configuration of a sapphire sphere and Si plate, both covered with Cr and Au layers. Equation (4) has never been used in previous literature (including [1,2,19]) to obtain constraints on the parameters of the Yukawa-type interaction from the measurements of the Casimir force. When dealing with the Yukawa interaction for the experimental parameters of [1,2], the plate can be considered as infinitely large. This was proved in [20] and confirmed in [8]. The exact expression for F_{sp}^{Yu} acting between a homogeneous sphere and a plane plate of sufficiently large area and thickness (as in experiment [1,2]) was found in Eq. (6) of [21]. Applying this equation to the layer structure of [1,2], one arrives at

$$\begin{aligned} F_{sp}^{\text{Yu}}(a) = & -4\pi^2 G\alpha\lambda^3 e^{-a/\lambda} \left[\rho_{\text{Au}} \Phi(R, \lambda) \right. \\ & - (\rho_{\text{Au}} - \rho_{\text{Cr}}) \Phi(R - \Delta_{\text{Au}}^{(s)}, \lambda) e^{-\Delta_{\text{Au}}^{(s)}/\lambda} \\ & - (\rho_{\text{Cr}} - \rho_s) \Phi(R - \Delta_{\text{Au}}^{(s)} - \Delta_{\text{Cr}}^{(s)}, \lambda) \\ & \quad \times e^{-(\Delta_{\text{Au}}^{(s)} + \Delta_{\text{Cr}}^{(s)})/\lambda} \Big] \\ & \times \left[\rho_{\text{Au}} - (\rho_{\text{Au}} - \rho_{\text{Cr}}) e^{-\Delta_{\text{Au}}^{(p)}/\lambda} \right. \\ & \quad - (\rho_{\text{Cr}} - \rho_{\text{Si}}) e^{-(\Delta_{\text{Au}}^{(p)} + \Delta_{\text{Cr}}^{(p)})/\lambda} \\ & \quad \left. - \rho_{\text{Si}} e^{-(\Delta_{\text{Au}}^{(p)} + \Delta_{\text{Cr}}^{(p)} + \Delta_{\text{Si}}^{(p)})/\lambda} \right], \end{aligned} \quad (5)$$

where

$$\Phi(r, \lambda) \equiv \Phi^{\text{exact}}(r, \lambda) = r - \lambda + (r + \lambda) e^{-2r/\lambda}. \quad (6)$$

Note that if we put

$$\Phi(r, \lambda) = \Phi^{\text{PFA}}(r, \lambda) \equiv R, \quad (7)$$

Eq. (5) results in $F_{sp}^{\text{Yu, PFA}}(a)$ which is also expressible as

$$F_{sp}^{\text{Yu, PFA}}(a) = 2\pi R E^{\text{Yu}}(a), \quad (8)$$

where $E^{\text{Yu}}(a)$ is the Yukawa energy per unit area of two parallel plates covered with thin layers as described above. Note that Eq. (8) is nothing but the simplified formulation of the PFA. The negative derivative of $E^{\text{Yu}}(a)$ with respect to a results in the Yukawa pressure between two parallel plates in Eq. (3).

Note that the thickness of the lower plate mentioned in the explanations to Eq. (8) might be finite whereas the upper plate is a semispace (see paper [6]). This is because the force $F_{sp}^{\text{Yu,PFA}}(a)$ in the left-hand side of Eq. (8) can depend only on the three geometrical parameters: separation a , sphere radius R and thickness of the lower plate D_1 (if it is not much larger than a , as in the experiments under discussion). However, Eq. (6) of [8] which is analogous to our Eq. (8) (the use of P_{Yu} instead of E^{Yu} is a misprint) contains in the right-hand side the energy per unit area of two parallel plates where the upper plate is of some arbitrary finite thickness D_2 . This equation is incorrect because its left-hand side does not depend on D_2 . Using this incorrect formulation of the PFA, paper [8] performed an extensive investigation of the dependence of $F_{sp}^{\text{Yu,PFA}}$ on D_2 [see Eq. (9) and Figs. 2 and 3] which is physically meaningless.

We next compare the constraints on the Yukawa parameters α, λ obtained in [1,2] from Eqs. (2), (3) using the simplified PFA, and the exact results following from Eqs. (4)–(6). According to [8], the larger λ is, the greater is the error in the constraints obtained using the PFA. The strongest constraints following from the data of Refs. [1,2] were obtained within the interaction range $20 \text{ nm} < \lambda < 86 \text{ nm}$. Hence as the first example we compare the resulting constraints at $\lambda = 86 \text{ nm}$. For this fixed λ the strongest constraints follow at $a = 250 \text{ nm}$ where the half-width of the confidence interval is $\tilde{\Xi}(a) = 1.52 \text{ mPa}$ [1]. From the exact Eqs. (4)–(6) we then obtain

$$\alpha^{\text{exact}} = 2.88167 \times 10^{13} \approx 2.88 \times 10^{13}. \quad (9)$$

Using the simplified PFA for the Yukawa interaction i.e., from Eqs. (2) and (3), it follows that

$$\alpha^{\text{PFA}} = 2.88011 \times 10^{13} \approx 2.88 \times 10^{13}. \quad (10)$$

Note that only the latter value was obtained in [1], and used to plot the line separating the allowed and prohibited regions in the (α, λ) -plane. It follows from a comparison of Eqs. (9) and (10) that the constraints on α, λ obtained in [1] are exact, to the quoted level of precision.

As a second example, we consider the much larger value, $\lambda = 400 \text{ nm}$, in the beginning of the region where the constraints obtained from the torsion pendulum experiment are the most stringent [20]. For this λ the strongest constraints are obtained from the Casimir force data at $a = 400 \text{ nm}$ and the half-width of the confidence interval is $\tilde{\Xi}(a) = 0.45 \text{ mPa}$ [1]. In this case the exact results in Eqs. (4)–(6) yield

$$\alpha^{\text{exact}} = 2.03189 \times 10^{11} \approx 2.0 \times 10^{11}, \quad (11)$$

whereas Eqs. (2) and (3) using the simplified PFA for the Yukawa force lead to

$$\alpha^{\text{PFA}} = 2.02708 \times 10^{11} \approx 2.0 \times 10^{11}. \quad (12)$$

The latter value was used in [1] to plot the line separating the allowed and prohibited regions in the (α, λ) -plane.

Note that the half-width of the confidence interval $\tilde{\Xi}(a)$ is found from the total theoretical and experimental errors in the Casimir pressure determined at a 95% confidence level. It takes into account all constituent errors, including the random and systematic experimental errors. Also included are theoretical errors due to uncertainty of the optical data, and due to the use of the PFA. As a quantity obtained from the errors, $\tilde{\Xi}(a)$ is calculated to only two, or at maximum to three, significant figures [the latter happens at shortest separations alone where $\tilde{\Xi}(a)$ is relatively large]. It would thus be an evidently inconsistent result if the two different determinations of the constraints on (α, λ) with two theoretical expressions for $F_{sp}^{\text{Yu}}(a)$ differing by only 0.1% would lead to markedly different strengths of the resulting constraints (to say nothing of the constraints differing by 100%, as discussed in [8]).

One can conclude that within the whole range of λ considered in [1,2] the strength of the derived constraints is precisely the same irrespective of whether or not the simplified PFA was used for the calculation of the Yukawa-type interaction between the sphere and the plate.

The opposite conclusion mentioned above was arrived at in [8]. According to [8], exact computations of the Casimir force “for the recent Casimir sphere-plane experiment [21] (here [2])... gives a deviation from PFA of the order of 0.1% at the smallest value of $a/R \approx 0.001$ reached in the experiment ($a_{\text{min}} \approx 160 \text{ nm}$). Since the limits to non-Newtonian forces are obtained using the residuals in the Casimir theory-experiment comparison, in order to meaningfully replace the exact formula of the Yukawa force with its PFA approximation, the level of accuracy between these two should be therefore a small fraction, for instance, 10% of the accuracy with which the Casimir force is controlled by using PFA rather than the exact expression for the sphere-plane Casimir force. If this condition is not fulfilled, the derived limits could be off also by a large order of 100% correction. However, targeting a 10% accuracy level with respect to the Casimir theory-experiment accuracy implies deviations from $\eta = 1$ of 0.01%, which can be obtained... only in the range of λ below 100 nm. The presence of substrates with different densities...” leads to the situation that η “in the case of the experiment reported in [21] (here [2]), is equal to 1.00126, i.e. a correction already equal to 0.126%.” Thus, according to [8], the deviation of the quantity η from unity in the experiment [2] exceeds allowed deviations by a factor of 13 and this may lead to order of 100% correction to the obtained constraints.

This statement is based on a simple misunderstanding. The key point is that the relative error in the maximum allowed value of α obtained using the simplified PFA is determined by the sum of relative errors in $\tilde{\Xi}(a)$ and in the application of the PFA to calculate the Yukawa force. Keeping in mind that $\tilde{\Xi}(a)$ has a meaning of an absolute error, it is determined with only two or three significant figures independently of its value. From this it follows that $\tilde{\Xi}(a)$ can only be known with the relative error of about

0.5%. As can be seen from the above citations, [8] mistakenly links a demand to the value of the relative error introduced by the application of the PFA to calculate the Yukawa force (of about 0.01%) with the magnitude of $\tilde{\Xi}(a)$ (or its constituent part due to the application of the PFA to calculate the Casimir force). Actually, to obtain constraints on α with the relative error of about 1% from the data of any high precision future experiment on the measurement of the Casimir force, it is quite sufficient to calculate the Yukawa force with a relative error of about 0.1% [22]. For example, at $a = \lambda = 400$ nm this calculation can be safely performed using the simplified formulation of the PFA if the application conditions of this formulation ($a, \lambda \ll R, D_1$) are satisfied [6]. Thus, the conclusion of [8] that the limits derived in [1, 2] could have “a large order of 100% correction” is invalid.

Acknowledgements

E.F. was supported in part by DOE under Grant No. DE-76ER071428. G.L.K. and V.M.M. were supported by Deutsche Forschungsgemeinschaft, Grant No. GE 696/10–1. G.L.K. was also supported by the Grant of the Russian Ministry of Education P–184.

References

1. R.S. Decca, D. López, E. Fischbach, G.L. Klimchitskaya, D.E. Krause, V.M. Mostepanenko, *Eur. Phys. J. C* **51**, 963 (2007)
2. R.S. Decca, D. López, E. Fischbach, G.L. Klimchitskaya, D.E. Krause, V.M. Mostepanenko, *Phys. Rev. D* **75**, 077101 (2007)
3. J. Blocki, J. Randrup, W.J. Swiatecki, C.F. Tsang, *Ann. Phys. (N.Y.)* **105**, 427 (1997)
4. E. Fischbach, C.L. Talmadge, *The Search for Non-Newtonian Gravity* (Springer, New York, 1999)
5. M. Bordag, G.L. Klimchitskaya, U. Mohideen, V.M. Mostepanenko, *Advances in the Casimir Effect* (Oxford University Press, Oxford, 2009)
6. R.S. Decca, E. Fischbach, G.L. Klimchitskaya, D.E. Krause, D. López, V.M. Mostepanenko, *Phys. Rev. D* **79**, 124021 (2009)
7. B.V. Derjaguin, *Kolloid. Z.* **69**, 155 (1934)
8. D.A.R. Dalvit, R. Onofrio, *Phys. Rev. D* **80**, 064025 (2009)
9. H. Gies, K. Klingmüller, *Phys. Rev. Lett.* **96**, 220401 (2006)
10. M. Bordag, V. Nikolaev, *J. Phys. A: Math. Theor.* **41**, 164002 (2008)
11. P.A. Maia Neto, A. Lambrecht, S. Reynaud, *Phys. Rev. A* **78**, 012115 (2008)
12. A. Canaguier-Durand, P.A. Maia Neto, I. Cervero-Pelaez, A. Lambrecht, S. Reynaud, *Phys. Rev. Lett.* **102**, 230404 (2009)
13. A. Wirzba, *J. Phys. A: Math. Theor.* **41**, 164003 (2008)
14. D.E. Krause, R.S. Decca, D. López, E. Fischbach, *Phys. Rev. Lett.* **98**, 050403 (2007)
15. T. Emig, R. L. Jaffe, *J. Phys. A: Math. Theor.* **41**, 164001 (2008)
16. T. Emig, *J. Stat. Mech.* P04007 (2008)
17. Particle Data Group, *Phys. Lett. B* **667**, 1 (2008)
18. G.L. Klimchitskaya, U. Mohideen, V.M. Mostepanenko, *Rev. Mod. Phys.* **81**, 1827 (2009)
19. R.S. Decca, D. López, E. Fischbach, G.L. Klimchitskaya, D.E. Krause, V.M. Mostepanenko, *Ann. Phys. (N.Y.)* **318**, 37 (2005)
20. M. Bordag, B. Geyer, G.L. Klimchitskaya, V.M. Mostepanenko, *Phys. Rev. D* **58**, 075003 (1998)
21. M. Bordag, B. Geyer, G.L. Klimchitskaya, V.M. Mostepanenko, *Phys. Rev. D* **62**, 011701(R) (2000)
22. V.B. Bezerra, G.L. Klimchitskaya, V.M. Mostepanenko, C. Romero, *Phys. Rev. D* **81**, 055003 (2010)